



Coastal Fisheries Creel Report Card

2021

VAITUPU

Introduction

This Coastal Fisheries Creel Report Card summarises the results of monitoring key indicators during creel surveys being carried out by Tuvalu Fisheries Department.

The Key indicators we use to show the health of the resources and state of overfishing are:

Indicator 1: Percentage of fishes that are landed which are smaller than the size at which at least 50% of the fish can breed (called length at maturity, L_m). This value should decline and approach zero as management actions improve, followed by improvements in the fisheries resources.

This is an indicator of **overfishing**.

Indicator 2: Catch of fishes per unit of effort (CPUE). We use the weight (kg) of fishes being landed: (a) per fisher per hour spent fishing and (b) per fishing trip. The values for Indicator 2 should increase as things improve. That is, fishers should be able to catch more fish in less time.

This is an indicator of **abundance** of the fishery as well as the **efficiency** of the fishing method.

Results

Overall status of Vaitupu's coastal resources is poor, with an average of 55% of the fishes caught being undersized. This is well above the national average of 35%.

The ideal % of fishes being landed that are undersized is 0, so any actions that will reduce this to lower levels is a step in the right direction and is expected to lead to improvements in the resources.

IDEAL: % UNDERSIZED should DECLINE over time and approach 0%




Figure 1: Percentage of fishes being landed undersized by year +/-SE. The sample size (n) is reported in blue.

Green arrow = good trend

Red arrow = bad trend

There was a significant increase in the percentage of undersized fish landed in Vaitupu between 2016 and 2019 (although there is not much data for 2019). This trend was reversed in 2021, which is a good signal.

Every fish should have the chance to breed at least once to ensure the resources can be replenished.

For Indicator 2, the total weight of fish being landed per fisher per hour spent fishing shows a slight decline for most fishing methods between 2017 and 2018. The CPUE for trolling appears to have increased in 2019, but is based on only 2 creel surveys (Figure 2).




Figure 2: Indicator 2a. Weight (in kg) of fishes landed per fisher per hour spent fishing across Tuvalu 2015-2021. There was no method data available for 2020.




Figure 3: Indicator 2b. (in kg) of fishes landed per fisher per fishing trip across Tuvalu 2015-2021.

The weight of fishes landed per fisher per entire fishing trip as Indicator 2b show similar trends to Indicator 2a – there is a slight decline between 2017 and 2018 for all fishing methods, and an increase in trolling in 2019 (Figure 3).

Catch per unit of effort (CPUE) should INCREASE over time in a well-managed fishery.

Conclusions

Overall, there has been little improvement to the health of coastal fisheries since surveys begun. The data suggest that between 2017 and 2018 there was more effort being used to catch fish – a greater number of which were undersized. More consistent data is needed to better understand these trends.

Management plans need to be developed and implemented more efficiently to improve the health of Tuvalu’s coastal fisheries.

Note: The catch reported do not include offshore fish species such as Atu (skipjack tuna). These pelagic species accounted for 22% of the species landed that were recorded in the creel surveys (2015-2021).

Why are some figures different from the previous report card?

This is due to a number of reasons:

1. We have received more data from the years 2015-2019
2. We have more accurate information on size of maturity from recently published studies
3. We have now included size of maturity data for 30 extra species
4. We have displayed CPUE by fishing method

Appendix I: Size of maturity (L_m) for top 50 species

Table 1 is part of indicator 1. It shows the breakdown of species that have 50% or more fishes landed that are undersized. A value of 100 means that all fishes landed are undersized. The ideal value for a well-managed fishery is 0. Blank cells indicate that no catch has been recorded for that species in that year. This table shows that many of the species being monitored are being caught undersized, and this varies by year.

The species are listed in order of their abundance in the catch landed (% of total catch).

Table 1: List of species for which size at maturity (L_m) is known, showing percentages landed which are undersized (2015-2021)

| | Scientific Name | Local Name | % of total catch | 2015 | 2016 | 2017 | 2018 | 2019 | 2021 |
|----|------------------------------------|--------------------------------|------------------|------|------|------|------|------|------|
| 1 | <i>Acanthurus triostegus</i> | Manini, Koinava | 15.5% | | 24 | 40 | 74 | | 14 |
| 2 | <i>Crenimugil crenilabis</i> | Kanase | 6.7% | 100 | | 59 | | | |
| 3 | <i>Liza vaigiensis</i> | Kafakafa | 4.9% | | | 84 | | | 100 |
| 4 | <i>Lutjanus fulvus</i> | Tagau, Takape | 2.1% | | | 29 | | | 31 |
| 5 | <i>Epinephelus macrospilos</i> | Gatala (Ff), fÄpuku (Nm) | 1.8% | | 100 | 100 | 100 | 100 | |
| 6 | <i>Kyphosus vaigiensis</i> | Nanue (Ff, Nm) | 1.3% | | | 87 | | | |
| 7 | <i>Lutjanus monostigma</i> | Taiva | 1.1% | | 0 | 53 | | | 100 |
| 8 | <i>Epinephelus merra</i> | Gatalaliki | 1.1% | | | 21 | | | 100 |
| 9 | <i>Epinephelus maculatus</i> | Fapuku | 0.6% | | | 100 | | | |
| 10 | <i>Acanthurus lineatus</i> | Ponelolo, Alogo, Pone hamao | 0.6% | | 0 | 29 | 80 | | |
| 11 | <i>Parupeneus barberinus</i> | Malili, Kaivete | 0.5% | | | 7 | | | |
| 12 | <i>Anyperodon leucogrammicus</i> | Gatala lautalo, Gatala lautala | 0.5% | | | 100 | | | |
| 13 | <i>Naso lituratus</i> | Maninilakau | 0.5% | | | 0 | | | |
| 14 | <i>Myripristis violacea</i> | Malau | 0.4% | | | 0 | | | 0 |
| 15 | <i>Lethrinus obsoletus</i> | Tanutanu | 0.4% | | | 44 | | | 20 |
| 16 | <i>Plectropomus areolatus</i> | Tonu gatala | 0.3% | | | 89 | | | |
| 17 | <i>Caranx sexfasciatus</i> | Teu | 0.3% | | 100 | 85 | | | 100 |
| 18 | <i>Ctenochaetus binotatus</i> | Pone, uli | 0.3% | | 17 | 22 | | | |
| 19 | <i>Parupeneus cyclostomus</i> | Kaivete piniki | 0.2% | | | 22 | | | |
| 20 | <i>Mulloidichthys vanicolensis</i> | Kalo | 0.2% | | | 0 | | | |